


## Chapter 10.6

# Smooth & Cardiac Muscles

---

Cardiac muscle



Smooth muscle



# Smooth Muscle

---

- Some smooth muscles are able to contract without nerves
- Other smooth muscle cells receive nerve stimulation from autonomic nervous system (Note: not somatic motor fibers as in skeletal muscle)
- SM capable of mitosis and hyperplasia
- Injured smooth muscle regenerates well

# Smooth Muscle

---

- composed of myocytes that have a **fusiform shape**
- one nucleus, located near the middle of the cell
- **no visible striations**
  - reason for the name 'smooth muscle'
  - thick and thin filaments are present, but not aligned with each other
- z discs are absent and replaced by **dense bodies**
  - well ordered array of protein masses in cytoplasm
  - protein plaques on the inner face of the plasma membrane
- cytoplasm contains extensive cytoskeleton of intermediate filament
  - attach to the membrane plaques and dense bodies
  - provide mechanical linkages between the thin myofilaments and the plasma membrane
- sarcoplasmic reticulum is scanty and there are no T tubules
- $\text{Ca}^{2+}$  needed for muscle contraction comes from the ECF by way of  $\text{Ca}^{2+}$  channels in the sarcolemma

# Stimulation of Smooth Muscle


---

- smooth muscle is **involuntary**
- can may contract without nervous stimulation
  - can contract in response to chemical stimuli
    - hormones, carbon dioxide, low pH, and oxygen deficiency
    - in response to stretch
    - single unit smooth muscle in stomach and intestines has **pacemaker cells** that set off waves of contraction throughout the entire layer of muscle
- most smooth muscle is innervated by **autonomic nerve fibers**
  - can trigger and modify contractions
  - **stimulate smooth muscle with either acetylcholine or norepinephrine**
  - can have contrasting effects
    - relax the smooth muscle of arteries
    - contract smooth muscles of the bronchioles

# 2 Types of Smooth Muscle


- **multiunit smooth muscle**

- occurs in some of the largest arteries and pulmonary air passages, in piloerector muscles of hair follicle, and in the iris of the eye
- ANS (autonomic nervous system) innervation similar to skeletal muscle
  - terminal branches of a nerve fiber synapse with individual myocytes and form a **motor unit**
  - each motor unit contracts independently of the others




(a) Multiunit smooth muscle

## 2 Types of Smooth Muscle




- **single-unit smooth muscle**
  - more widespread
  - occurs in most blood vessels, in the digestive, respiratory, urinary, and reproductive tracts
  - also called visceral muscle
    - often in two layers
    - inner circular
    - outer longitudinal
  - myocytes of this cell type are electrically coupled to each other by **gap junctions**
  - they directly stimulate each other and a large number of cells contract as a single unit

## 2 Types of Smooth Muscle




- in single unit smooth, each autonomic nerve fibers has up to 20,000 beadlike swelling called **varicosities**
  - each contains synaptic vesicles and a few mitochondria
  - nerve fiber passes amid several myocytes and stimulates all of them at once when it releases its neurotransmitter
- no motor end plates, but receptors scattered throughout the surface
- diffuse junctions – no one-to-one relationship between nerve fiber and myocyte

# Stimulation of Smooth Muscle



# Layers of Visceral Muscle

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



# Smooth Muscle Contraction and Relaxation

---

- contraction is triggered by  $\text{Ca}^{+2}$ , energized by ATP, and achieved by sliding thin past thick filaments
- contraction begins in response to  $\text{Ca}^{+2}$  that enters the cell from ECF, a little internally from sarcoplasmic reticulum
  - $\text{Ca}^{+2}$  channels open to allow  $\text{Ca}^{+2}$  to enter cell
  - voltage, ligand, and mechanically-gated (stretching)
- calcium binds to **calmodulin** on thick filaments
  - activates **myosin light-chain kinase** – adds phosphate to regulatory protein on myosin head
  - **myosin ATPase**
    - hydrolyzing ATP
    - enables myosin similar power and recovery strokes like skeletal muscle
  - thick filaments pull on thin ones - thin ones pull on dense bodies and membrane plaques
    - force is transferred to plasma membrane and entire cell shortens
    - puckers and twists like someone wringing out a wet towel

# Contraction and Relaxation

---


- very slow in comparison to skeletal muscle
  - latent period in skeletal 2 msec, smooth muscle 50 - 100 msec
  - tension peaks at about 500 msec (0.5 sec)
  - declines over a period of 1 – 2 seconds
  - slows myosin ATPase enzyme and slow pumps that remove  $\text{Ca}^{+2}$
  - $\text{Ca}^{+2}$  binds to calmodulin instead of troponin // activates kinases and ATPases that hydrolyze ATP
  - smooth muscle makes most of its ATP aerobically

# Contraction and Relaxation

---

- smooth muscle resistant to fatigue
  - **latch-bridge mechanism** - heads of myosin molecules do not detach from actin immediately
  - do not consume more ATP immediately
  - maintains tetanus tonic contraction (**smooth muscle tone**)
    - arteries – vasomotor tone
    - intestinal tone

# Contraction of Smooth Muscle



# Stretching Smooth Muscle

---

- **stretch** can open mechanically-gated calcium channels in the sarcolemma causing contraction
  - **peristalsis** – waves of contraction brought about by food distending the esophagus or feces distending the colon // propels contents along the organ
- **stress-relaxation response** (receptive relaxation)
  - helps hollow organs gradually fill (urinary bladder)
  - when stretched, tissue briefly contracts then relaxes – helps prevent emptying while filling

# Contraction and Stretching

---

- smooth muscle **contracts forcefully even when greatly stretched**
  - allows hollow organs such as the stomach and bladder to fill and then expel their contents efficiently
  - smooth muscle can be anywhere from half to twice its resting length and still contract powerfully
  - skeletal muscle cannot contract forcefully if overstretched

# Contraction and Stretching

---

- three reasons why
  - there are no z discs, so thick filaments cannot butt against them and stop contraction
  - since the thick and thin filaments are not arranged in orderly sarcomeres, stretching does not cause a situation where there is too little overlap for cross-bridges to form
  - the **thick filaments of smooth muscle have myosin heads along their entire length**, so cross-bridges can form anywhere
- **plasticity** – the ability to adjust its tension to the degree of stretch // a hollow organ such as the bladder can be greatly stretched yet not become flabby when it is empty

# Cardiac Muscle




# Cardiac Muscle

---

- limited to the heart where it functions to pump blood
- required properties of cardiac muscle
  - contraction with regular rhythm
  - muscle cells of each chamber must contract in unison
  - contractions must last long enough to expel blood
  - must work in sleep or wakefulness, with out fail, and without conscious attention
  - must be highly resistant to fatigue

## Cardiac muscle



# Cardiac Muscle

---

- characteristics of cardiac muscle cells
  - **striated** like skeletal muscle, but myocytes (cardiocytes) are **shorter and thicker**
  - each myocyte is joined to several others at the uneven, notched linkages – **intercalated discs**
    - appear as thick dark lines in stained tissue sections
    - electrical **gap junctions** allow each myocyte to directly stimulate its neighbors
    - mechanical junctions that keep the myocytes from pulling apart
  - **sarcoplasmic reticulum less developed**, but T tubules are larger and admit supplemental  $\text{Ca}^{2+}$  from the extracellular fluid
  - damaged cardiac muscle cells repair by **fibrosis**
    - a **little mitosis** observed following heart attacks
    - not in significant amounts to regenerate functional muscle

# Cardiac Muscle

- can contract without need for nervous stimulation
  - contains a **built-in pacemaker** that rhythmically sets off a wave of electrical excitation
  - wave travels through the muscle and triggers contraction of heart chambers
  - **autorhythmic** – because of its ability to contract rhythmically and independently
- **autonomic nervous system** does send nerve fibers to the heart
  - can increase or decrease heart rate and contraction strength
- very slow twitches - does not exhibit quick twitches like skeletal muscle
  - maintains tension for about 200 to 250 msec
  - gives the heart time to expel blood
- uses **aerobic respiration** almost exclusively
  - rich in myoglobin and glycogen
  - has especially **large mitochondria**
    - 25% of volume of cardiac muscle cell
    - 2% of skeletal muscle cell with smaller mitochondria
- very adaptable with respect to fuel used
- very vulnerable to interruptions of oxygen supply
- **highly fatigue resistant**