

Chapter 21 (1)



Blood Vessels and Blood Circulation



The human body has 60,000 miles of blood vessels.

This is nearly enough blood vessels to circle the earth 2 ½ times.

When you gain one pound of fat, you add another mile of blood vessels.




Lecture Objectives



- **Compare and contrast the structure of an artery, arteriole, vein, venule, and capillary**
- **Discuss the structure and function of a capillary network**


Blood Vessels

1 mm




- arteries - carry blood away from heart
- veins - carry blood back to heart
- capillaries - connect smallest arteries to smallest veins
- capillaries allow for the exchange of fluid, ions, & small molecules between the blood and the interstitial space
- arteries and veins have **three tunics** (layers)






(c) Capillary




(a) Artery



Tunica Interna (tunica intima)


- Cells that lines the blood vessel
- **endothelium** – simple squamous epithelium overlying a basement membrane and a sparse layer of loose connective tissue
 - function as a **selectively permeable barrier**
 - normally **repels blood cells and platelets** that may adhere to it and form a clot (endothelium secrete **prostacyclin – make surface “slippery”**)
 - when tissue around vessel is inflamed, the endothelial cells **produce cell-adhesion molecules** that induce **leukocytes to adhere to the surface**
 - allows leukocytes to congregate in tissues where their defensive actions are needed



Tunica Media



- middle layer
- consists of smooth muscle and connective tissue with collagen and elastic fibers
- strengthens vessel and prevents blood pressure from rupturing vessels
- **vasomotion** – changes in diameter of the blood vessel brought about by smooth muscle action /// regulated by vasomotion center in medula oblongata




Note: at end of arterial system are small diameter vessels called arterioles // the vasomotor center is able to control blood pressure by regulating smooth muscle contraction --- this is known as peripheral resistance.

The volume of blood on the arterial side also contributes to the blood pressure.

Tunica Externa

- outermost layer
- consists of loose connective tissue
- merges with that of neighboring blood vessels, nerves, or other organs
- anchors the vessel
- provides passage for small nerves, lymphatic vessels
- **vasa vasorum** – small vessels that supply blood to at least the outer half of the blood vessels
- blood from the lumen is thought to nourish the inner half of the blood vessel by diffusion



Arteries

- Carry blood away from the heart
- Maybe classified by their function
 - Elastic arteries (largest – garden hose size /// also function as a pressure reservoirs)
 - Distributing arteries (medium size sometimes referred to as muscular / pencil to string size)
 - Resistance arterioles (small arteries – able to regulate blood flow into capillary beds and change blood pressure)

Distributing Arteries

Distributing arteries are also called conducting arteries

- **these arteries have “names”**
- arteries distal to pulmonary trunk and aorta // e.g. common carotid, subclavian, and common iliac arteries
- have a layer of elastic tissue, **internal elastic lamina**, at the border between intima and media
- **external elastic lamina** at the border between media and externa
- expand during systole, recoil during diastole // this lessens fluctuations in blood pressure

Elastic Arteries


Walls stretched as blood received by pulmonary trunk and aorta

Stretching vessel walls “stores energy” because of elastic component of the vessel wall


Elastic recoil in pulmonary trunk and aorta keep blood flowing while ventricles are in diastole

Resistance Arteries


- Why are arteries called resistance vessels // because they have thick smooth muscle walls
 - Resilient wall structure that can resist dilation due to high blood pressure
 - Arterioles (small arteries just before capillaries) are “regulated” by the vasomotor center
 - Two functions of the **vasomotor center** (located in medulla oblongata)
 - **Regulate blood pressure & regulate blood distribution**



(a) Elastic aorta and arteries stretch during ventricular contraction



(b) Elastic aorta and arteries recoil during ventricular relaxation



Cross-sectional area

Velocity

Aorta

Arteries

Arterioles

Capillaries

Venules

Veins

Venae cavae

Arterial Sense Organs



- Sensory structures in the walls of the **aorta and carotid arteries** monitor blood pressure and blood chemistry
- *Transmit information to brain stem to regulate*
 - 1) *heart rate* *
 - 2) *vasomotion (vasomotor center)* *
 - 3) *respiration*
- *Location of sense organs*
 - *carotid baroreceptor (also called carotid sinus)*
 - *aortic baroreceptor (also called aortic sinus)*
 - *carotid chemoreceptor*
 - *aortic chemoreceptor*

 - *Please Note: the chemoreceptors are more important in regulating respiration.*


Arterial Sense Organs

- **aortic & carotid sinuses** // baroreceptors (pressure sensors)
 - **most important in regulation of heart**
 - in walls of internal carotid artery & aortic arch
 - monitors blood pressure – signaling brainstem
 - decreased heart rate and vessels dilation in response to high blood pressure

- **aortic & carotid bodies** // chemoreceptors
 - oval bodies near branch of common carotids
 - monitor blood chemistry
 - **mainly transmit signals to the brainstem respiratory centers**
 - adjust respiratory rate to stabilize pH, CO₂, and O₂
 - one to three in walls of aortic arch
 - same function as carotid bodies



Capillaries



- Site where nutrients, wastes, and hormones pass between the blood and tissue fluid through the walls of the vessels (exchange vessels)

- the ‘business end’ of the cardiovascular system
- composed of **endothelium and basal lamina**
- Capillaries are absent or scarce in tendons, ligaments, epithelia, cornea and lens of the eye

- **Three Capillary Types**

- distinguished by ease with which substances pass through their walls
- structural differences that account for their greater or lesser permeability


- **Continuous Capillaries**
- **Fenestrated Capillaries**
- **Sinusoid Capillaries**




Continuous Capillaries

- occur in most tissues
- **endothelial cells** have **tight junctions** forming a continuous tube with **intercellular clefts**
 - allow passage of solutes such as glucose
- **pericytes** wrap around the capillaries and contain the same contractile protein as muscle
 - contract and regulate blood flow



Continuous Capillary






(a) Continuous capillary formed by endothelial cells

Fenestrated Capillary



b: Courtesy of S. McNutt




(b) Fenestrated capillary




Fenestrated Capillaries

- common in kidneys and small intestine
- organs that require rapid absorption or filtration
- endothelial cells riddled with holes called **filtration pores (fenestrations)**
 - spanned by very thin glycoprotein layer
 - allows passage of only small molecules

Sinusoid in Liver





(c) Sinusoid



Sinusoids

- discontinuous capillaries
- liver, bone marrow, spleen
- irregular blood-filled spaces with large fenestrations
- allow proteins (albumin), clotting factors, and new blood cells to enter the circulation

Capillary Beds




- Capillaries organized into networks called **capillary beds**
- Situated between an arteriole and venule
- A passageway connects the arteriole to the venule // two segments
 - **Metarteriole** – proximal to arteriole / no smooth muscle but smooth muscle at junction between metarteriole and capillary
 - **precapillary sphincters** control which beds are well perfused
 - when sphincters open - capillaries are well perfused with blood and engage in exchanges with the tissue fluid
 - when sphincters closed - blood bypasses the capillaries and flows through thoroughfare channel to venule
 - **Thoroughfare channel** – point beyond the metarteriole that continues through capillary bed to venule

Arterioles and Metarterioles

- **Arterioles**

- Responsible for “**peripheral resistance**”
- smallest arteries
- control amount of blood flowing into various organs
- thicker tunica media in proportion to their lumen than large arteries
- very little tunica externa

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.




Arterioles and Metarterioles

- **Metarterioles**

- short vessels that link arterioles to capillaries
- muscle cells form a **precapillary sphincter** about entrance to capillary
 - constriction of these sphincters reduces or **shuts off blood flow through their respective capillaries**
 - diverts blood to other tissues


Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



(a) Sphincters open

Capillary Bed Sphincters Open

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



(a) Sphincters open

approximately one billion capillaries

no cell more than 40 to 80 micrometers from capillary


four to six cell widths

Pre-capillary sphincters are not controlled by vasomotor center

Pre-capillary sphincters regulated by “local regulation”


when sphincters are open, the capillaries are well perfuse ///
three-fourths of the capillaries of the body are normally shut down

Capillary Bed Sphincters Closed




(b) Sphincters closed

when the sphincters are closed, little to no blood flow into capillary bed
(e.g. when skeletal muscles at rest)



(b) Sphincters contracted: blood flowing through thoroughfare channel



(a) Sphincters relaxed: blood flowing through capillaries


Autoregulation



- **Autoregulation = Local Control** = the ability of the tissues to regulate its own blood supply
 - **Metabolic Theory of Auto-Regulation**
 - This principle applies to the capillary beds
 - If tissue is not adequately perfused with blood then carbon dioxide accumulate and stimulates vasodilation which increases perfusion
 - Bloodstream delivers oxygen and removes carbon dioxide
 - When carbon dioxide levels drop then smooth muscle constricts
 - Why does this make sense? Explain.


Capillary Filtration and Reabsorption

- capillary **filtration** at arterial end
- capillary **reabsorption** at venous end
- Variations within organ type
 - Location
 - Glomeruli of kidney
 - devoted to filtration
 - Alveolar capillary of lungs
 - devoted to absorption
 - activity or trauma
 - increases filtration






Capillary Filtration and Reabsorption




Net "6 mm Hg" out. **Where does the fluid go?**

Structure of a Capillary Bed with Lymphatic Capillaries and Their Afferent Vessels



How much fluid is not recovered at the end of the capillary bed?



Key:
 BHP = Blood hydrostatic pressure
 IFHP = Interstitial fluid hydrostatic pressure
 BCOP = Blood colloid osmotic pressure
 IFOP = Interstitial fluid osmotic pressure
 NFP = Net filtration pressure

Net filtration at arterial end of capillaries (20 liters per day) Net reabsorption at venous end of capillaries (17 liters per day)


Net filtration pressure (NFP) = (BHP + IFOP) - (BCOP + IFHP)
 Pressures promoting filtration Pressures promoting reabsorption


Arterial end
NFP = (35 + 1) - (26 + 0) = 10 mmHg
Result: Net filtration

Venous end
NFP = (16 + 1) - (26 + 0) = -9 mmHg
Result: Net reabsorption

Veins (The Capacitance Vessels)

- greater capacity (volume) for blood containment than arteries
- thinner walls, flaccid, less muscular and elastic tissue
- collapse when empty, expand easily
- have steady blood flow
- merge to form larger veins
- subjected to relatively low blood pressure /// remains 10 mm Hg with little fluctuation





Venules & Veins

- **Venules** – smallest veins, occur just past the capillary bed
 - even more porous than capillaries // also exchange fluid with surrounding tissues
 - tunica interna with a few fibroblasts and no muscle fibers
 - most leukocytes emigrate from the bloodstream through venule walls
- **Muscular venules** – up to 1 mm in diameter
 - 1 or 2 layers of smooth muscle in tunica media /// have a thin tunica externa

Venules & Veins

- **medium veins** – up to 10 mm in diameter
 - thin tunica media and thick tunica externa
 - tunica interna in this area forms **venous valves**
 -
- **large veins** – larger than 10 mm
 - some smooth muscle in all three tunics
 - thin tunica media with moderate amount of smooth muscle
 - tunica externa is thickest layer
 - contains longitudinal bundles of smooth muscle
 - Examples /// venae cavae, pulmonary veins, internal jugular veins, and renal veins


Sinuses

- Venous sinuses
 - veins with extremely thin walls
 - large lumen, and no smooth muscle
 - dural venous sinus in the brain
 - coronary sinus in the heart
 - These vessels are not capable of vasomotion
 - Function as blood reservoirs

Mechanisms of Venous Return




- **Venous return** – the flow of blood back to the heart
 - pressure gradient
 - blood pressure is the most important force in venous return
 - 7-13 mm Hg venous pressure towards heart
 - venules (12-18 mm Hg) to **central venous pressure** – point where the venae cavae enter the heart (~5 mm Hg)
 - **skeletal muscle pump** – occurs in the limbs
 - contracting muscle compressed vein between muscles to move blood towards heart
 - deep veins in legs have one way valves
 - veins in arm also have valves, but fewer than in legs



Skeletal Muscle Pump

Propels venous blood back toward the heart



(a) Contracted skeletal muscles

(b) Relaxed skeletal muscles

Other Mechanisms of Venous Return

- gravity drains blood from head and neck
- thoracic (respiratory) pump
 - inhalation - thoracic cavity expands and thoracic pressure decreases, abdominal pressure increases forcing blood upward
/// central venous pressure fluctuates
 - 2mm Hg- inhalation, 6mm Hg-exhalation
 - blood flows faster with inhalation
- “cardiac suction” caused by expanding atria

Venous Return and Physical Activity



- exercise increases venous return in many ways
 - heart beats faster, harder increasing CO and BP
 - vessels of skeletal muscles, lungs, and heart dilate and increase flow
 - increased respiratory rate, increased action of thoracic pump
 - increased skeletal muscle pump

Venous Return and Physical Activity

- venous pooling occurs with inactivity
 - venous pressure not great enough to force blood upward
 - with prolonged standing, may be low enough to cause dizziness or fainting
 - prevented by tensing leg muscles to activate skeletal muscle pump
 - jet pilots wear pressure suits


Varicose Veins




- blood pools in the lower legs in people who stand for long periods stretching the veins
 - cusps of the valves pull apart in enlarged superficial veins further weakening vessels
 - blood backflows and further distends the vessels, their walls grow weak and develop into **varicose veins**
- hereditary weakness, obesity, and pregnancy also promote problems
- **hemorrhoids** are varicose veins of the anal canal

Normal and Varicose Veins


Normal veins





Varicose veins




A Normal vein




B Varicose vein








Circulatory Routes




- Most common route
 - heart → arteries → arterioles → capillaries → venules → veins
 - passes through only **one network of capillaries** from the time it leaves the heart until the time it returns
- **Portal system** // blood flows through **two capillary** networks before returning to heart
 - artery → capillary → portal vein → capillary → vein
 - Three locations:
 - 1) between hypothalamus and anterior pituitary
 - 2) kidneys (peritubular capillaries)
 - 3) between intestines and liver



- **anastomosis** – the point where two blood vessels merge
- **arteriovenous anastomosis (shunt)**
 - artery flows directly into vein bypassing capillaries
- **venous anastomosis**
 - most common
 - one vein empties directly into another
 - reason vein blockage less serious than an arterial blockage
- **arterial anastomosis**
 - two arteries merge
 - provides collateral (alternative) routes of blood supply to a tissue
 - coronary circulation and around joints






Aneurysm




- **Weak point in an wall of an artery**
 - forms a thin-walled, bulging sac that pulsates with each heartbeat and may rupture at any time
 - **dissecting aneurysm** - blood accumulates between the tunics of the artery and separates them, usually because of degeneration of the tunica media
 - **most common sites** // abdominal aorta, renal arteries, and arterial circle at the base of the brain
 - can cause pain by putting pressure on other structures
 - can rupture causing hemorrhage
 - result from congenital weakness of the blood vessels or result of trauma or bacterial infections such as syphilis
 - most common cause is atherosclerosis and hypertension


Aneurysm



Saccular Aneurysm




Fusiform Aneurysm




Lecture Objectives Stop

Review of Blood Vessels



■ = Oxygenated blood
■ = Deoxygenated blood






Anterior view of coronary arteries and their major branches

SCHEME OF DISTRIBUTION






Superior vena cava
Right pulmonary artery
Pulmonary trunk
Right pulmonary veins
Right atrium
Right ventricle
Right lung
Inferior vena cava

Arch of aorta
Ascending aorta
Left pulmonary artery
Left pulmonary veins
Left atrium
Left ventricle
Left lung
Diaphragm
Abdominal aorta



(a) Anterior view

SCHEME OF DISTRIBUTION


Abdominal aorta




SCHEME OF DISTRIBUTION




SCHEME OF DISTRIBUTION




SCHEME OF DISTRIBUTION






(a) Anterior view


(b) Posterior view



Overall anterior view of the principal veins




Anterior view





Right lateral view



SCHEME OF DRAINAGE




SCHEME OF DRAINAGE




SCHEME OF DRAINAGE




SCHEME OF DRAINAGE




SCHEME OF DRAINAGE






(b) Anterior view of deep veins




(a) Anterior view of superficial veins





(a) Anterior view

(b) Posterior view




(c) Posterior view of superficial veins of hand







(c) Scheme of fetal circulation



(b) Scheme of principal blood vessels of hepatic portal circulation and arterial supply and venous drainage of liver



(a) Anterior view of veins draining into the hepatic portal vein



(b) Scheme of pulmonary circulation