

AP / Bio 2700
Chapter Study Guide Answers
Endocrinology C18

1. Cells of Exocrine glands deliver secretions onto a surface by way of a duct. // Endocrine glands transport their secretions (hormones) to target tissue by using the blood stream.
2. Nervous System = electrochemical communication = fast (milliseconds) // Endocrine System = chemical messengers (hormones) = slow (minutes – hours – days – months)
3. A small number of enzymes can produce a lot of product since they are not consumed in the catalytic reaction and can be used over and over // Hormones direct changes in the target tissue's metabolism which is regulated by enzymes.
4. Chemical messenger
5. Hormones influence cellular metabolism throughout the body because the hormones are transported throughout the body by the blood // Paracrines are also chemical messengers secreted by cells however paracrines do not enter the blood stream and only affect cells in the immediate area of their production.
6. Hormone produced by variety of cells types // secreted into blood // only cells throughout body with receptor matched to hormone will bind hormone // only after hormone binds to target tissue with appropriate receptor will hormone modify metabolism of target tissue
7. Hypophyseal pituitary axis shows the regulatory relationship between the hypothalamus and pituitary. Hypothalamus produces chemical messengers (hormones) with travel down infundibulum to stimulate release of hormones from pituitary (i.e. the “master gland” of the endocrine system // most influential endocrine gland in human body)
8. Glucagon, growth hormone, epinephrine, norepinephrine, cortisol, corticosterone
9. Insulin
10. (see separate worksheet for list of hormones)
11. Tract tells you that this structure consists of axons. They connect areas of the hypothalamus that produces hormones (ADH and oxytocin) with the posterior pituitary that release these two hormones
12. A portal system is a blood circulatory circuit which consists of two capillary beds between an artery and a vein. In this case it allows hormones to be produced in

- one area (hypothalamus) and transported directly to a specific location (anterior pituitary)
13. Hormone receptors may be either on the surface of the target cell's plasma membrane or inside the cell (e.g. the nucleoplasm)
 14. Blood glucose levels rise.
 15. Diabetes implies an increase in urine volume. In diabetes insipidus the cause for the increase in urine volume is due to the reduced secretion of ADH. // In diabetes mellitus type I the cause is lack of insulin secretion which causes high blood glucose levels. This then results in glucose exceeding the kidney's transport maximum for glucose, the kidney can not reabsorb all the glucose it filters and some glucose "spills" into the urine which increases urine volume by an osmotic force (i.e. osmotic drag). // In DM type II blood glucose levels are elevated because the target tissues are insensitive to insulin. The kidney responds to elevated glucose levels similar to what was experienced in type I.
 16. Parathyroid hormone stimulates the breakdown of bone calcium-phosphate crystals therefore it allows calcium to move into the blood // Calcitonin stimulates new bone growth which requires moving calcium from the blood into the bone (i.e. lowers blood calcium levels)
 17. Aldosterone and cortisol (much lower concentrations androgens and estrogens // not principle secretory gland for these two hormones)
 18. Hydrophilic hormones dissolve in blood // hydrophobic hormones need to be "transported" because do not dissolve in blood – often hydrophobic hormones are attached to albumin which is a soluble blood protein.
 19. Adrenal gland.
 20. See figures in book.
 21. See separate worksheet.
 22. Oxytocin regulation is our reference model for positive feedback. It is a "neuron – endocrine pathway" seen in both the let down of milk from the mammary gland and in parturition. Nerve tracts go into hypothalamus to increase production-release of oxytocin into blood which goes to target tissue where it further stimulates the nerve tracts going to the hypothalamus // TSH is our reference model for negative feedback. TSH is secreted from the pituitary and its target tissue is the thyroid glands which secrete thyroid hormone which then goes back to the pituitary and stops the secretion of TSH (i.e. negative feedback)

23. Steroids (e.g. estrogen and testosterone) // peptides (e.g. insulin) // monoamines (e.g. epinephrine)
24. Steroids are hydrophobic and require a carrier molecule in the blood which has the effect of making the steroid hormones have longer half lives. Peptides and monoamines are soluble in blood and tend to have shorter half lives.
25. Second messengers are also called metabotropic. These are transmembrane proteins with a receptor on the outer face of the plasma membrane which the hormone may engage. If the hormone fits into the receptor then on the inner face of the transmembrane protein a "G protein" is released which activates a different transmembrane protein which "makes" a new molecule which is the "second messenger". // peptide and monoamine hormones must use second messenger systems to change metabolism of their target tissues.
26. Ovaries secrete estrogen (stimulated by FSH). The ovaries corpus luteum secretes progesterone and inhibin.
27. Testes secrete testosterone and inhibin. FSH and interstitial cell hormone regulate male gonadal tissue hormonal secretions.